

Towards the AdS dual of SYK

Vladimir Rosenhaus
KITP

D. Gross, V.R, “All-point correlation functions in SYK”,
1710.08113

May 4, 2018

In this talk I will make some comments on the status of the AdS dual of SYK

In this talk I will make some comments on the status of the AdS dual of SYK

I will briefly describe the results for the computation of all large N correlation functions in SYK.

In this talk I will make some comments on the status of the AdS dual of SYK

I will briefly describe the results for the computation of all large N correlation functions in SYK.

This, in principal, fully determines the bulk Lagrangian: all the masses and couplings of the (infinite) tower of bulk fields.

In this talk I will make some comments on the status of the AdS dual of SYK

I will briefly describe the results for the computation of all large N correlation functions in SYK.

This, in principal, fully determines the bulk Lagrangian: all the masses and couplings of the (infinite) tower of bulk fields.

However, we are far from a clear physical understanding of what the bulk theory actually is.

I will first remind you of some basic aspects of the canonical example of AdS/CFT: the duality between string theory in $\text{AdS}_5 \times S^5$ and $\mathcal{N}=4$ super Yang-Mills.

In $\mathcal{N}=4$ there are six fundamental scalars (which are $N \times N$ matrices), out of which one can build many single-trace operators, such as $\text{Tr}(XXYXXY)$.

In $\mathcal{N}=4$ there are six fundamental scalars (which are $N \times N$ matrices), out of which one can build many single-trace operators, such as $\text{Tr}(XXYXXY)$.

One can compute the dimensions of these operators. At large 't Hooft coupling, most of these operators acquire a large dimension, as one would expect.

Each of these single-trace operators $\text{Tr}(XXYXXY)$ is dual to a different field in the bulk, whose mass is set by the dimension of the corresponding operator.

Each of these single-trace operators $\text{Tr}(XXYXXY)$ is dual to a different field in the bulk, whose mass is set by the dimension of the corresponding operator.

At large 't Hooft coupling, the stringy modes acquire a large mass. If it were not for the Kaluza-Klein modes on the S^5 , there would only be a few light fields.

Each of these single-trace operators $\text{Tr}(XXYXXY)$ is dual to a different field in the bulk, whose mass is set by the dimension of the corresponding operator.

At large 't Hooft coupling, the stringy modes acquire a large mass. If it were not for the Kaluza-Klein modes on the S^5 , there would only be a few light fields.

The 't Hooft coupling is dual to the ratio of the AdS scale to the string scale, so this is what we would expect.

In SYK, the fundamental fields are fermions χ_i out of which we can construct $O(N)$ singlets,

$$\mathcal{O}_n \sim \frac{1}{N} \sum_{i=1}^N \chi_i \partial_\tau^{1+2n} \chi_i$$

In SYK, the fundamental fields are fermions χ_i out of which we can construct $O(N)$ singlets,

$$\mathcal{O}_n \sim \frac{1}{N} \sum_{i=1}^N \chi_i \partial_\tau^{1+2n} \chi_i$$

We will say that each of these is dual to a scalar ϕ_n in AdS_2

In SYK, the fundamental fields are fermions χ_i out of which we can construct $O(N)$ singlets,

$$\mathcal{O}_n \sim \frac{1}{N} \sum_{i=1}^N \chi_i \partial_\tau^{1+2n} \chi_i$$

We will say that each of these is dual to a scalar ϕ_n in AdS_2

From the SYK four-point function, we know the dimensions h of these operators, at strong coupling. They differ by an order-one amount from $2n+1$.

In SYK, the fundamental fields are fermions χ_i out of which we can construct $O(N)$ singlets,

$$\mathcal{O}_n \sim \frac{1}{N} \sum_{i=1}^N \chi_i \partial_\tau^{1+2n} \chi_i$$

We will say that each of these is dual to a scalar ϕ_n in AdS_2

From the SYK four-point function, we know the dimensions h of these operators, at strong coupling. They differ by an order-one amount from $2n+1$.

The lowest dimension operator is $h=2$, and is special. It is responsible for the breaking of conformal invariance and maximal chaos, and is dual to the dilaton. It is roughly like the graviton (talks by Maldacena, Verlinde).

The dual of SYK therefore has a tower of fields in the bulk, whose masses are roughly spaced by even integers.

The dual of SYK therefore has a tower of fields in the bulk, whose masses are roughly spaced by even integers.

This is like string theory, with an AdS scale of order the string scale.

The dual of SYK therefore has a tower of fields in the bulk, whose masses are roughly spaced by even integers.

This is like string theory, with an AdS scale of order the string scale.

But this of course is not the usual string theory, since there are far fewer fields.

The bulk action is of the form

$$S_{bulk} = \int d^2x \sqrt{g} \left[\frac{1}{2} (\partial \phi_n)^2 + \frac{1}{2} m_n^2 \phi_n^2 + \frac{1}{\sqrt{N}} \lambda_{nmk} \phi_n \phi_m \phi_k + \dots \right]$$

We have already discussed the masses.


The bulk action is of the form

$$S_{bulk} = \int d^2x \sqrt{g} \left[\frac{1}{2} (\partial \phi_n)^2 + \frac{1}{2} m_n^2 \phi_n^2 + \frac{1}{\sqrt{N}} \lambda_{nmk} \phi_n \phi_m \phi_k + \dots \right]$$


We have already discussed the masses.

To find the cubic couplings, we must compute the SYK three-point function of bilinears (six-point function of fermions), for the quartic couplings we need the four-point function of bilinears, and so on.


I will now make some technical remarks about the computation of all-point correlation functions in SYK



=




+




$$C_{h_1 h_2 h_3}$$

$$\mathcal{O} \sim \chi_i \partial_\tau^{1+2n} \chi_i$$




$$c_{h_1 h_2 h_3}$$


=




+



$$\mathcal{O} \sim \chi_i \partial_\tau^{1+2n} \chi_i$$



$$\int_{\mathcal{C}} \frac{dh}{2\pi i} \tilde{\rho}(h) c_{h_1 h_2 h} c_{h_3 h_4 h} \mathcal{F}_{h_i}^h(\tau_i)$$




$$\int_{\mathcal{C}} \frac{dh}{2\pi i} \rho(h) \mathcal{F}_{\Delta}^h(\tau_i) \quad \rho(h) \sim \frac{k(h)}{1 - k(h)}$$



- h-space is to the conformal group, $SL(2, \mathbb{R})$, what Fourier space is to translations

$$\int \frac{dp}{2\pi} f(p) e^{ipx}$$


Maldacena Stanford
 Polchinski, V.R
 Kitaev



$$\int_{\mathcal{C}} \frac{dh}{2\pi i} \tilde{\rho}(h) c_{h_1 h_2 h} c_{h_3 h_4 h} \mathcal{F}_{h_i}^h(\tau_i)$$




$$\int_{\mathcal{C}} \frac{dh_a}{2\pi i} \tilde{\rho}(h_a) \int_{\mathcal{C}} \frac{dh_b}{2\pi i} \tilde{\rho}(h_b) c_{h_1 h_2 h_a} c_{h_a h_3 h_b} c_{h_b h_4 h_5} \mathcal{F}_{h_i}^{h_a, h_b}(\tau_i)$$



$$\int_{\mathcal{C}} \frac{dh}{2\pi i} \tilde{\rho}(h) c_{h_1 h_2 h} c_{h_3 h_4 h} \mathcal{F}_{h_i}^h(\tau_i)$$

- These are simple rules for summing an infinite number of diagrams. It doesn't matter that the four-point function is made up of ladders. These apply to any four-point functions.



$$\int_C \frac{dh}{2\pi i} \tilde{\rho}(h) c_{h_1 h_2 h} c_{h_3 h_4 h} \mathcal{F}_{h_i}^h(\tau_i)$$

- These are simple rules for summing an infinite number of diagrams. It doesn't matter that the four-point function is made up of ladders. These apply to any four-point functions.
- This is not just an OPE expansion. The $c_{h_1 h_2 h_3}$ are the analytically extended OPE coefficients of the single-trace operators. The four-point function is a sum of conformal blocks of single-trace operators and double-trace operators. This emerges upon closing the contour. Somehow, the analytically extended single-trace OPE coefficients encode the information about the double-trace OPE coefficients.

The only result for the correlation function that I will use in the rest of the talk is the three-point function of bilinears.

The bilinear three-point function is composed of the hypergeometric function ${}_4F_3$ and gamma functions. It simplifies in the limit of large dimensions. From the three-point function, it is trivial to get the bulk cubic couplings,

The bilinear three-point function is composed of the hypergeometric function ${}_4F_3$ and gamma functions. It simplifies in the limit of large dimensions. From the three-point function, it is trivial to get the bulk cubic couplings,

$$\lambda_{n_1 n_2 n_3} \approx \frac{N!}{\Gamma(N - 2n_1 + \frac{1}{2})\Gamma(N - 2n_2 + \frac{1}{2})\Gamma(N - 2n_3 + \frac{1}{2})}$$

$$N = n_1 + n_2 + n_3 \qquad n_i \gg 1$$

The bilinear three-point function is composed of the hypergeometric function ${}_4F_3$ and gamma functions. It simplifies in the limit of large dimensions. From the three-point function, it is trivial to get the bulk cubic couplings,

$$\lambda_{n_1 n_2 n_3} \approx \frac{N!}{\Gamma(N - 2n_1 + \frac{1}{2})\Gamma(N - 2n_2 + \frac{1}{2})\Gamma(N - 2n_3 + \frac{1}{2})}$$

$$N = n_1 + n_2 + n_3 \qquad n_i \gg 1$$

Notice that if one sets $n_1=n_2=n_3$, this grows exponentially.

$$S_{bulk} = \int d^2x \sqrt{g} \left[\frac{1}{2} (\partial \phi_n)^2 + \frac{1}{2} m_n^2 \phi_n^2 + \frac{1}{\sqrt{N}} \lambda_{nmk} \phi_n \phi_m \phi_k + \dots \right]$$

Summary: The bulk has a tower of fields, with masses roughly spaced by even integers, and cubic couplings of the form on the previous slide.

$$S_{bulk} = \int d^2x \sqrt{g} \left[\frac{1}{2} (\partial \phi_n)^2 + \frac{1}{2} m_n^2 \phi_n^2 + \frac{1}{\sqrt{N}} \lambda_{nmk} \phi_n \phi_m \phi_k + \dots \right]$$

Summary: The bulk has a tower of fields, with masses roughly spaced by even integers, and cubic couplings of the form on the previous slide.

The spectrum looks roughly like a Kaluza-Klein reduction of a single scalar field in $AdS_2 \times S^1$ (talk by Jevicki for a precise match)

$$S_{bulk} = \int d^2x \sqrt{g} \left[\frac{1}{2} (\partial \phi_n)^2 + \frac{1}{2} m_n^2 \phi_n^2 + \frac{1}{\sqrt{N}} \lambda_{nmk} \phi_n \phi_m \phi_k + \dots \right]$$

Summary: The bulk has a tower of fields, with masses roughly spaced by even integers, and cubic couplings of the form on the previous slide.

The spectrum looks roughly like a Kaluza-Klein reduction of a single scalar field in $AdS_2 \times S^1$ (talk by Jevicki for a precise match)

However, the cubic couplings seem qualitatively different from what one gets from a Kaluza-Klein reduction.

In fact, the cubic couplings are, heuristically, of the type one gets in string theory.

In fact, the cubic couplings are, heuristically, of the type one gets in string theory.

For string theory in 10 dimensions, one has creation operators to excite the string modes in the 8 transverse directions: a_n^1, \dots, a_n^8

In fact, the cubic couplings are, heuristically, of the type one gets in string theory.

For string theory in 10 dimensions, one has creation operators to excite the string modes in the 8 transverse directions: a_n^1, \dots, a_n^8

The cubic couplings are roughly, $\langle a_n a_m^\dagger a_k^\dagger \rangle$

In fact, the cubic couplings are, heuristically, of the type one gets in string theory.

For string theory in 10 dimensions, one has creation operators to excite the string modes in the 8 transverse directions: a_n^1, \dots, a_n^8

The cubic couplings are roughly, $\langle a_n a_m^\dagger a_k^\dagger \rangle$

This gives some combinatorial factors, vaguely like what we found.


To make an actual statement, we need string theory in 2 dimensions, not 10. However, in 2 dimensions there are no transverse directions.

To make an actual statement, we need string theory in 2 dimensions, not 10. However, in 2 dimensions there are no transverse directions.

One can try to instead allow longitudinal motion: a string that is expanding and contracting.

Bardeen, Bars, Hanson, Peccei, '76

Maldacena, '05, Maldacena Stanford, '16




To make an actual statement, we need string theory in 2 dimensions, not 10. However, in 2 dimensions there are no transverse directions.

One can try to instead allow longitudinal motion: a string that is expanding and contracting.

Bardeen, Bars, Hanson, Peccei, '76

Maldacena, '05, Maldacena Stanford, '16

Semiclassically, the spectrum agrees with SYK



To make an actual statement, we need string theory in 2 dimensions, not 10. However, in 2 dimensions there are no transverse directions.


One can try to instead allow longitudinal motion: a string that is expanding and contracting.

Bardeen, Bars, Hanson, Peccei, '76

Maldacena, '05, Maldacena Stanford, '16

Semiclassically, the spectrum agrees with SYK

To get the cubic couplings, we would need to quantize the string. It is unclear how to do this.



A famous solvable two-dimensional field theory is the 't Hooft model: large N QCD with fermions.

A famous solvable two-dimensional field theory is the 't Hooft model: large N QCD with fermions.

The spectrum of meson masses in the 't Hooft model is like that of the string with longitudinal motion: $m_n^2 \sim n$

A famous solvable two-dimensional field theory is the 't Hooft model: large N QCD with fermions.

The spectrum of meson masses in the 't Hooft model is like that of the string with longitudinal motion: $m_n^2 \sim n$

Let us therefore study the 't Hooft model, placed in AdS. Is this the dual of SYK?

A famous solvable two-dimensional field theory is the 't Hooft model: large N QCD with fermions.

The spectrum of meson masses in the 't Hooft model is like that of the string with longitudinal motion: $m_n^2 \sim n$

Let us therefore study the 't Hooft model, placed in AdS. Is this the dual of SYK?

It is unlikely that it would precisely match the dual of SYK, but it is conceivable that it is close

A famous solvable two-dimensional field theory is the 't Hooft model: large N QCD with fermions.

The spectrum of meson masses in the 't Hooft model is like that of the string with longitudinal motion: $m_n^2 \sim n$

Let us therefore study the 't Hooft model, placed in AdS. Is this the dual of SYK?

It is unlikely that it would precisely match the dual of SYK, but it is conceivable that it is close

We have not yet succeeded in solving the 't Hooft model in AdS, so I do not know the answer.

Summary


- SYK has a tower of fermion bilinear, $O(N)$ singlets
 $\mathcal{O}_n \sim \chi_i \partial_\tau^{1+2n} \chi_i$ with anomalous dimensions of order 1.

Summary

- SYK has a tower of fermion bilinear, $O(N)$ singlets
 $\mathcal{O}_n \sim \chi_i \partial_\tau^{1+2n} \chi_i$ with anomalous dimensions of order 1.
 - We have computed all-point correlation functions of these operators.
-

Summary

- SYK has a tower of fermion bilinear, $O(N)$ singlets
 $\mathcal{O}_n \sim \chi_i \partial_\tau^{1+2n} \chi_i$ with anomalous dimensions of order 1.
- We have computed all-point correlation functions of these operators.
- The correlation functions take a simple form, when written correctly.



$$\int_{\mathcal{C}} \frac{dh}{2\pi i} \tilde{\rho}(h) c_{h_1 h_2 h} c_{h_3 h_4 h} \mathcal{F}_{h_i}^h(\tau_i)$$

- The \mathcal{O}_n are assumed to be dual to scalars ϕ_n in AdS.

- The \mathcal{O}_n are assumed to be dual to scalars ϕ_n in AdS.
- Their dimensions determine the masses of the scalars, and their three-point function determines the cubic couplings.

- The \mathcal{O}_n are assumed to be dual to scalars ϕ_n in AdS.
- Their dimensions determine the masses of the scalars, and their three-point function determines the cubic couplings.
- The cubic couplings, of the large mass fields, take a combinatorial form,

$$\lambda_{n_1 n_2 n_3} \approx \frac{N!}{\Gamma(N - 2n_1 + \frac{1}{2})\Gamma(N - 2n_2 + \frac{1}{2})\Gamma(N - 2n_3 + \frac{1}{2})}$$

$$N = n_1 + n_2 + n_3 \qquad n_i \gg 1$$

- The form of the cubic couplings perhaps suggests a stringy origin.

- The form of the cubic couplings perhaps suggests a stringy origin.
- Some possibilities are a string with longitudinal motion, or 2d QCD with fermions. We do not yet know if this gives the correct cubic couplings.

- It is possible that one should be trying to find the dual not of SYK, but of tensor models. Tensor models have the same melonic Feynman diagrams as SYK, but their symmetry is $O(N) \times O(N) \times O(N)$,

$$S = \int d\tau \left(\frac{1}{2} \psi_{abc} \partial_\tau \psi_{abc} + \frac{1}{4} g \psi_{abc} \psi_{ade} \psi_{fbe} \psi_{fdc} \right)$$

Gurau; Witten;
Klebanov & Tarnopolsky

- It is possible that one should be trying to find the dual not of SYK, but of tensor models. Tensor models have the same melonic Feynman diagrams as SYK, but their symmetry is $O(N) \times O(N) \times O(N)$,

$$S = \int d\tau \left(\frac{1}{2} \psi_{abc} \partial_\tau \psi_{abc} + \frac{1}{4} g \psi_{abc} \psi_{ade} \psi_{fbe} \psi_{fdc} \right)$$

Gurau; Witten;
Klebanov & Tarnopolsky

- There are a huge number of singlets. This gives an extraordinary large number of bulk fields, far far more than SYK, and far more than string theory.

- It is possible that one should be trying to find the dual not of SYK, but of tensor models. Tensor models have the same melonic Feynman diagrams as SYK, but their symmetry is $O(N) \times O(N) \times O(N)$,

$$S = \int d\tau \left(\frac{1}{2} \psi_{abc} \partial_\tau \psi_{abc} + \frac{1}{4} g \psi_{abc} \psi_{ade} \psi_{fbe} \psi_{fdc} \right)$$

Gurau; Witten;
Klebanov & Tarnopolsky

- There are a huge number of singlets. This gives an extraordinary large number of bulk fields, far far more than SYK, and far more than string theory.
- What the dual of tensor models might be is a mystery.